

Kate O'Hanlan, M. D.
F. A. C. O. G., S. G. O., F. A. C. S.
Gynecologic Oncology Associates
4370 Alpine Road, Suite 104
Portola Valley, CA 94028-7927

Phone: (650)-851-6669

ohanlan.com

FAX: (650) 851-9747

Brief review of recent evidence about supracervical hysterectomy

According to data from 26,000 women in the Women's Health Initiative, similar rates of cystocele, rectocele, and stress and urge incontinence were observed in women with and without hysterectomy.¹ Prolapse was only predicted by obesity and parity.

Neither total or subtotal hysterectomy affect urinary continence or sexual function.²⁻⁶

Complication rates are not lower for supracervical hysterectomy than total hysterectomy.⁵⁻⁷

Pap smears must be continued per standard indications as HPV is common. Some suggest that HPV testing be done before subtotal hysterectomy,⁸ especially for younger women, or women who are not in monogamous relationships. Cervical cancer rates continue to rise throughout a woman's life with an age specific incidence of 30/100,000 at age 80, more than double that of age 30 to 40.

Problems can develop from the residual cervical stump, such as pain^{7,9}, prolapse³, recurrence of fibroids¹⁰ or development of cancer¹¹⁻¹⁵, ectopic pregnancy¹⁶, tubo-ovarian abscess¹⁷, necrosis and shock¹⁸ and may ultimately require resection in 25%^{9,19}.

Up to 25% continue to have cyclic bleeding,^{7,20} This cyclic bleeding is evidence of persistent responsive cervical glandular tissue.

Unopposed estrogen (ERT) has been associated with development of cervical adenocarcinoma.²¹ Furthermore, obesity, common among 67% of American women, is a known risk factor for cervical adenocarcinoma, just as it is for endometrial carcinoma²²

Thus, patients who have had supracervical hysterectomy who have significant menopausal symptoms must take combined estrogen+progestin (HRT) rather than estrogen only (ERT) to minimize risk of cervical cancer.^{21,22} However, the Women's Health Initiative shows that HRT, and not ERT, increases risk of breast cancer, clots, stroke and heart disease.²³ When the entire uterus is removed, ERT is all that is needed for menopausal symptoms.

1. Hendrix SL, Clark A, Nygaard I, Aragaki A, Barnabei V, McTiernan A. Pelvic organ prolapse in the Women's Health Initiative: gravity and gravidity. *Am J Obstet Gynecol.* Jun 2002;186(6):1160-1166.
2. Kim DH, Lee YS, Lee ES. Alteration of sexual function after classic intrafascial supracervical hysterectomy and total hysterectomy. *J Am Assoc Gynecol Laparosc.* Feb 2003;10(1):60-64.
3. Thakar R, Ayers S, Clarkson P, Stanton S, Manyonda I. Outcomes after total versus subtotal abdominal hysterectomy. *N Engl J Med.* Oct 24 2002;347(17):1318-1325.
4. Kuppermann M, Summitt RL, Jr., Varner RE, et al. Sexual functioning after total compared with supracervical hysterectomy: a randomized trial. *Obstet Gynecol.* Jun 2005;105(6):1309-1318.
5. Learman LA, Summitt RL, Jr., Varner RE, et al. A randomized comparison of total or supracervical hysterectomy: surgical complications and clinical outcomes. *Obstet Gynecol.* Sep 2003;102(3):453-462.
6. Scott JR, Sharp HT, Dodson MK, Norton PA, Warner HR. Subtotal hysterectomy in modern gynecology: a decision analysis. *Am J Obstet Gynecol.* 1997;176(6):1186-1191; discussion 1191-1182.
7. van der Stege JG, van Beek JJ. Problems related to the cervical stump at follow-up in laparoscopic supracervical hysterectomy. *Jsls.* 1999;3(1):5-7.
8. Ford JF, Feinstein SM. Human papillomavirus testing before elective supracervical hysterectomy. *J Low Genit Tract Dis.* Oct 2005;9(4):230-231.
9. Nezhat CH, Nezhat A, Borhan S. Laparoscopic Removal of the Cervical Stump after Supracervical Hysterectomy for Persistent Pelvic Pain and Endometriosis. *J Am Assoc Gynecol Laparosc.* 1996;3(4, Supplement):S34.
10. Bojahr B, Lober R, Romer T, Schwesinger G. [Large cervix myoma after supracervical hysterectomy]. *Zentralbl Gynakol.* 1996;118(8):468-470.
11. Mendez LE, Penalver M, McCreath W, Bejarano P, Angioli R. Radical vaginal trachelectomy after supracervical hysterectomy. *Gynecol Oncol.* Jun 2002;85(3):545-547.
12. Schneider A. Recurrence of unclassifiable uterine cancer after modified laparoscopic hysterectomy with morcellation. *Am J Obstet Gynecol.* Aug 1997;177(2):478-479.
13. Hannoun-Levi JM, Peiffert D, Hoffstetter S, Luporsi E, Bey P, Pernot M. Carcinoma of the cervical stump: retrospective analysis of 77 cases. *Radiother Oncol.* May 1997;43(2):147-153.
14. Waxman M, Waxman JS, Alinovi V. Heterologous malignant mixed Mullerian tumor of the cervical stump. *Gynecol Oncol.* Dec 1983;16(3):422-428.
15. Markowska J, Markowska A. [Total or subtotal hysterectomy?]. *Ginekol Pol.* 2000;71(1):34-38.
16. Pasic R, Scobee J, Tolar B. Ectopic pregnancy months after laparoscopic supracervical hysterectomy. *J Am Assoc Gynecol Laparosc.* Feb 2004;11(1):94-95.
17. Tohya T, Yoshimura T, Onoda C. Tubo-ovarian abscess occurring 16 years after supracervical hysterectomy. *Infect Dis Obstet Gynecol.* 2003;11(3):167-169.

18. Huang JY, Ziegler C, Tulandi T. Cervical stump necrosis and septic shock after laparoscopic supracervical hysterectomy. *J Minim Invasive Gynecol.* Mar-Apr 2005;12(2):162-164.
19. Okaro EO, Jones KD, Sutton C. Long term outcome following laparoscopic supracervical hysterectomy. *Bjog.* 2001;108(10):1017-1020.
20. Ghomi A, Hantes J, Lotze EC. Incidence of cyclical bleeding after laparoscopic supracervical hysterectomy. *J Minim Invasive Gynecol.* May-Jun 2005;12(3):201-205.
21. Lacey JV, Jr., Brinton LA, Barnes WA, et al. Use of hormone replacement therapy and adenocarcinomas and squamous cell carcinomas of the uterine cervix. *Gynecol Oncol.* Apr 2000;77(1):149-154.
22. Lacey JV, Jr., Swanson CA, Brinton LA, et al. Obesity as a potential risk factor for adenocarcinomas and squamous cell carcinomas of the uterine cervix. *Cancer.* Aug 15 2003;98(4):814-821.
23. Rossouw JE, Anderson GL, Prentice RL, et al. Risks and benefits of estrogen plus progestin in healthy postmenopausal women: principal results From the Women's Health Initiative randomized controlled trial. *Jama.* Jul 17 2002;288(3):321-333.

Sexual function after hysterectomy and supracervical hysterectomy

Davis, A. R. (2000). "Recent advances in female sexual dysfunction." *Curr Psychiatry Rep* 2(3): 211-4.

Female sexuality has received little scientific study. Recently, increased interest in this field has generated new research in the epidemiology, pathophysiology, and pharmacotherapy of female sexual dysfunction (FSD). A new FSD classification system has been proposed. Although sexual difficulties are highly prevalent among women, the degree of associated distress is unknown. Risk factors for FSD are probably both psychologic and physiologic. Aging or menopause is associated with lubrication difficulties, which can be treated with hormone replacement.

Hysterectomy seems more likely to result in improvement rather than deterioration of sexual functioning. Depression may be a predictor of sexual dysfunction after hysterectomy. Vasoactive agents are currently being evaluated as treatment for female sexual arousal disorder. The most important advance in the study of female sexual function is the recent surge of interest in this relatively unexplored field.

Galyer, K. T., H. M. Conaglen, et al. (1999). "The effect of gynecological surgery on sexual desire." *J Sex Marital Ther* 25(2): 81-8.

This study compared the impact of surgery on sexual desire in women who had undergone hysterectomy with women whose nongynecological surgery was of a similar status. Women aged 30 to 65 completed mailed questionnaires including the Sexual Desire Questionnaire, Hurlbert Index of Sexual Desire, and the Sexual

Desire Inventory. Other areas assessed were affect, mood awareness, sexual anxiety, and androgen levels. **Analysis of the questionnaire measures showed no differences in sexual desire for gynecological versus nongynecological surgery, or across the different types of hysterectomy.** There was no relation between the level of androgens and the women's sexual desire levels. The implications of these findings are discussed with regard to the potential impact of surgery on sexuality, as well as future research in the area.

Rannestad, T., O. J. Eikeland, et al. (2001). "Are the physiologically and psychosocially based symptoms in women suffering from gynecological disorders alleviated by means of hysterectomy?" *J Womens Health Gend Based Med* 10(6): 579-87.

Women experiencing gynecological disorders can suffer from a diversity of symptoms and problems. To what extent women are relieved of their physiologically and psychosocially based symptoms by hysterectomy is an important issue. This study aims to evaluate the long-term impact of hysterectomy on such symptoms and to compare the findings with those of a control group. Women with gynecological disorders experienced physiologically based symptoms, such as pelvic pain and urinary incontinence, to a greater degree than did the control group ($p < 0.05$), whereas bowel function and menopausal symptoms were equally distributed in the two groups. Furthermore, these women experienced psychosocially based symptoms, such as fatigue and insomnia, to a greater degree than the control group ($p < 0.05$), whereas the groups did not differ in psychological well-being or sexual function. After hysterectomy, however, pelvic pain and sleeping disturbances were alleviated, and the state of energy and urinary function were improved ($p < 0.00$). Bowel function, menopausal symptoms, psychological well-being, sexual function, and self-perception as a woman were not influenced by the operation. Both 6 and 12 months after hysterectomy, no differences between the patients and the control group were found, except for the tendency of patients not to gain weight after the operation. **The symptoms are mostly improved or unchanged after hysterectomy, and additional problems do not seem to follow the operation.**

Rhodes, J. C., K. H. Kjerulff, et al. (1999). "Hysterectomy and sexual functioning." *Jama* 282(20): 1934-41.

CONTEXT: Women considering hysterectomy often are concerned about its potential effects on their sexual functioning but the effects of hysterectomy on sexual functioning remain unclear. OBJECTIVE: To examine changes in sexual functioning after hysterectomy. DESIGN AND SETTING: A 2-year prospective study (Maryland Women's Health Study) of hysterectomy, which included measures of sexual functioning prior to hysterectomy and at 6, 12, 18, and 24 months after hysterectomy, performed during 1992 and 1993. PATIENTS: Of

1299 women interviewed prior to hysterectomy, 1101 (84.8%) completed the study and provided information about their sexual functioning. Most were between the ages of 35 and 49 years, white, married or living with a partner, and high school graduates. MAIN OUTCOME MEASURES: Frequency of sexual relations, dyspareunia, orgasm, vaginal dryness, and sexual desire. RESULTS: The percentage of women who engaged in sexual relations increased significantly from 70.5% before hysterectomy to 77.6% and 76.7% at 12 and 24 months after hysterectomy. The rate of frequent dyspareunia dropped significantly from 18.6% before hysterectomy to 4.3 % and 3.6% at 12 and 24 months after hysterectomy. The rates of not experiencing orgasms dropped significantly from 7.6% before hysterectomy to 5.2% and 4.9% at 12 and 24 months after hysterectomy. Low libido rates also decreased significantly from 10.4% before hysterectomy to 6.3% and 6.2% at 12 and 24 months after hysterectomy. The distribution of women not reporting vaginal dryness in the past month improved significantly from 37.3% before hysterectomy to 46.8% and 46.7% at 12 and 24 months after hysterectomy. Prehysterectomy depression was associated with experiencing dyspareunia, vaginal dryness, low libido, and not experiencing orgasms after hysterectomy. CONCLUSIONS: **Sexual functioning improved overall after hysterectomy. The frequency of sexual activity increased and problems with sexual functioning decreased.**

Utian, W. H. and P. P. Boggs (1999). "The North American Menopause Society 1998 Menopause Survey. Part I: Postmenopausal women's perceptions about menopause and midlife." *Menopause* 6(2): 122-8.

OBJECTIVE: To collect information relevant to the mission of The North American Menopause Society (NAMS)--i.e., increasing understanding of menopause--by assessing perceptions held by postmenopausal women in the United States aged 50 to 65 years regarding their menopause transition and early postmenopausal years. DESIGN: During the period from June to July 1998, The Gallup Organization conducted 752 telephone interviews with a randomly selected sample of postmenopausal women aged 50 to 65 years from across the United States, based on questions developed by NAMS. In Part I of this survey, women were asked about their personal experiences with menopause, their health-related lifestyle changes since premenopause, their frequency of discussing menopause, and their rating of preparedness for menopause. Part II of this survey, including use of hormone replacement therapy as well as use of healthcare services, will be reported in a future communication from NAMS. RESULTS: The majority (51%) of the postmenopausal women surveyed reported being happiest and most fulfilled between the ages of 50 to 65 years, compared with when they were in their 20s (10%), 30s (17%), or 40s (16%). Many areas of their lives had improved since menopause, including family/home life, sense of personal fulfillment, ability to focus on hobbies or other interests, relationship with spouse/partner, and friendships. A majority (51%) said their sexual relationships had remained

unchanged. Approximately three-quarters of women surveyed reported making some type of health-related lifestyle change, such as stopping smoking, at menopause/midlife. Women who had undergone hysterectomy expressed more improvement than women with an intact uterus, especially in the areas of sexual relationships, spouse/partner relationships, personal fulfillment, and physical health; data are not available regarding the health state of these women before surgery or whether they experienced surgical menopause, but this improvement did not appear to be the result of hormone replacement therapy. Women tended to look to women from their own generation for menopause-related information and believed that they have prepared the younger generation for menopause better than they were prepared by their mothers' generation. Those surveyed advised younger women to engage in healthful activities and become knowledgeable so that they could make informed health decisions. **CONCLUSIONS:** Although the postmenopausal women surveyed had differing views of menopause as well as their perceptions of postmenopause compared with premenopause, the majority viewed menopause and midlife as the beginning of many positive changes in their lives and health. **Hysterectomy was a factor associated with improved sexual relationships, spouse/partner relationships, sense of personal fulfillment, and physical health.**

Weber, A. M., M. D. Walters, et al. (1999). "Functional outcomes and satisfaction after abdominal hysterectomy." *Am J Obstet Gynecol* 181(3): 530-5.

OBJECTIVES: Our aim was to compare urinary, lower gastrointestinal, and sexual function and to describe patients' expectations and satisfaction before and after hysterectomy. **STUDY DESIGN:** Forty-three women completed questionnaires before and about 1 year after abdominal hysterectomy for benign gynecologic conditions. Symptoms related to urinary, lower gastrointestinal, and sexual function and satisfaction with treatment were assessed. To account for multiple comparisons, only $P \leq .002$ was considered statistically significant. **RESULTS:** There were no statistically significant changes in urinary or bowel symptoms before and after hysterectomy, with preoperative symptoms resolving in some women after surgery and developing in others. Sexual function including orgasmic ability did not change in the 34 sexually active women. The level of satisfaction with treatment was very high. **CONCLUSIONS: Women experience high degrees of satisfaction and no consistent changes in urinary, lower gastrointestinal, or sexual function after abdominal hysterectomy for benign gynecologic conditions.**